



State-dependent atmospheric noise associated with ENSO

Jong-Seong Kug,^{1,2} Fei-Fei Jin,² K. P. Sooraj,¹ and In-Sik Kang¹

Received 21 September 2007; revised 6 November 2007; accepted 12 December 2007; published 1 March 2008.

[1] A strong relationship between ENSO (El Niño/Southern Oscillation) and atmospheric short-term variability in the near-surface zonal wind is reported in the present study. On one hand, anomalies in the variance of the short-term atmospheric variability over the western Pacific tend to lead El Niño development. On the other hand, the activity of the fast atmospheric variability over the central Pacific is simultaneously correlated to the NINO3.4 SST. The significant correlation exists over the broad range of atmospheric variability from the synoptic time scale to the inter-seasonal time scale. This finding supports the notion that the ENSO–state dependence of the atmospheric variability serves as a source of stochastic forcing for ENSO. Furthermore, it is demonstrated that there is a significant interdecadal change in this dependence. The state-dependent noise becomes much stronger in the recent period, which possibly explains the recent increase of the ENSO activities, consistent with the recent theory of noise-induced destabilization effect for ENSO. **Citation:** Kug, J.-S., F.-F. Jin, K. P. Sooraj, and I.-S. Kang (2008), State-dependent atmospheric noise associated with ENSO, *Geophys. Res. Lett.*, *35*, L05701, doi:10.1029/2007GL032017.

1. Introduction

[2] Since the largest El Niño and strongest activity of westerly wind events (WWE) were observed during the year 1997/98 ENSO event [McPhaden, 1999], a number of studies have suggested the importance of interaction between ENSO and relatively shorter timescale atmospheric variability such as the Madden-Julian Oscillation (MJO), and WWE [Moore and Kleeman, 1999; Kessler and Kleeman, 2000; Vecchi and Harrison, 2000; Zhang and Gottschalk, 2002; Yu et al., 2003; Kirtman et al., 2005; Wu and Kirtman, 2006]. Some of these studies showed that the enhanced MJO and WWE activities preceded the peak of El Niño by a few months [McPhaden, 1999; Harrison and Vecchi, 1997; Fink and Speth, 1997; Zhang and Gottschalk, 2002]. Others indicate that the MJO and WWEs are preceded by a large-scale SST anomaly [Luther et al., 1983; Gutzler, 1991; Hendon et al., 1999; Vecchi and Harrison, 2000; Kessler, 2001].

[3] Recent theoretical and modeling studies have elucidated the importance of this interaction between two different timescale phenomena [Lengaigne et al., 2004; Eisenman

et al., 2005; Perez et al., 2005; Zavala-Garay et al., 2005; Jin et al., 2007]. For instance, Jin et al. [2007] demonstrated in a conceptual framework that the interaction between ENSO and the fast atmospheric variability, as modeled by the state-dependent noise forcing or multiplicative noise forcing, enhances instability of ENSO, its ensemble spread, and generate ENSO asymmetry. They proposed a theory of noise-induced instability for ENSO through the interaction between ENSO and the fast atmospheric variability. Observational studies regarding the intensity and nature of the interaction between ENSO and the fast atmospheric variability state are needed in order to quantify the potential importance of the so-called state-dependent noise on ENSO. In this study, authors try to examine the interaction between ENSO and fast equatorial wind variability by using the 56-yr NCEP/NCAR reanalysis data and by focusing on the level and decadal variation of dependence of the atmospheric “noise” on the ENSO state.

2. Data and Methodology

[4] The data used in this study are monthly means of SST and daily 925-hPa zonal wind for the period of 1950–2005. The monthly mean SST data are from the improved extended reconstructed sea surface temperature version 2 (ERSST.v2) data set [Smith and Reynolds, 2004] created by the National Climate Data Center (NCDC); this data analysis uses monthly and 2° spatial superobservations, which are defined as individual observations averaged onto their 2° grid. The daily wind data are taken from the National Center for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR [Kalnay et al., 1996]). Further, we verified our results with the daily wind data sets from ERA40. Despite significant differences documented in other studies between aspects of the NCEP and ERA40 tropical Pacific wind stress [e.g., Wittenberg, 2004], we found that the results presented here are not sensitive to the reanalysis data sets. Daily anomalies are obtained after removing the climatologic annual cycle, which is obtained by averaging the all the daily data on the same calendar dates. A 2–180 day band pass with the LANCZOS filter (using 45 weights [Duchon, 1979]) is applied to the daily wind anomalies, in order to investigate relatively the fast atmospheric variability associated with ENSO. The filtered zonal wind anomaly data are averaged over 5°S–5°N latitudinal width, to only focus on the impact of atmospheric noise on the equatorial wave guide. Hereafter, the variability of the filtered wind is referred to as atmospheric “noise” for simplicity. Noted that our definition on the “noise” is somewhat limited because the noise occurs all space and time scales. In this study, we only focus on high-frequency atmospheric variability associated with

¹Climate Environment System Research Center, Seoul National University, Seoul, Korea.

²Department of Meteorology, School of Ocean and Earth Science and Technology, University of Hawaii at Manoa, Honolulu, Hawaii, USA.




Figure 1. Lag-regression (contour) and correlation (shading) coefficients of (a) SST (averaged over 5°S – 5°N) with respect to NINO3.4 SST, and (b) atmospheric noise variability (averaged over 5°S – 5°N). Lag (months) is indicated on the vertical axes, with negative values representing SST or atmospheric noise leading NINO3.4 SST. The unit of regression coefficients is dimensionless and $\text{m}^2 \text{s}^{-2} \text{K}^{-1}$, respectively.

ENSO. The variance of the filtered wind is calculated in 5-month moving window.

3. Results

[5] To examine the relation between ENSO and atmospheric noise variability, lag-regression coefficients are calculated with respect to the NINO3.4 SST, which is an anomalous SST averaged over 170 – 120°W , 5°S – 5°N . Noted that the statistical relation between ENSO and atmospheric variability is not sensitive to a choice of other ENSO indices such as NINO3 and NINO4 SSTs. Figure 1 shows the lag-regression coefficients for the recent 26-year period of 1980–2005. The SST anomalies are also displayed for comparison. In overall view, the enhanced noise variance seems to propagate eastward from the western Pacific at the El Niño onset phase to the eastern Pacific at the El Niño decaying period. In particular, stronger (weaker) activities for the atmospheric noise are observed over the western Pacific 7–10 months prior to the mature stage of El Niño (La Niña). The correlation coefficients are more than 0.5 with 99% confidence level. Note that the large tropical Pacific SST anomaly is not yet established by that time. The strong variance at El Niño onset phase is possibly linked to the eastward extension of warm pool [Kessler *et al.*, 2005; Eisenman *et al.*, 2005] and anomalous westerlies over the western Pacific [Seiki and Takayabu, 2007a, 2007b]. This indicates that the strong MJO and WWE activities play a role in triggering the development of El Niño, as previous studies have argued [McPhaden, 1999; Kessler and Kleeman, 2000; Boulanger *et al.*, 2001; Lengaigne *et al.*, 2004]. Therefore, the activity of the fast atmospheric variability can be used as a precursor for El Niño onset, one of the challenging problems in current El Niño prediction.

[6] A distinctive strong relationship between ENSO and atmospheric noise over the central Pacific is shown in Figure 1. The noise variance is simultaneously correlated with the NINO3.4 SST. The correlation coefficients are more than 0.6 with 99% confidence level. This significant

relationship was reported in previous studies focusing on MJO activity [Gutzler, 1991; Hendon *et al.*, 1999; Kessler, 2001; Tam and Lau, 2005] and WWEs [Luther *et al.*, 1983; Harrison and Vecchi, 1997; Seiki and Takayabu, 2007a]. The eastward extension of the warm pool [Kessler *et al.*, 1995] and background westerlies flow [Lau and Lau, 1992] can accompany the eastward shift of MJO events and WWEs during the El Niño mature phase. This close relationship between ENSO and noise variance supports the theoretical framework and this relation is modeled simply as idealized state-dependent noise [Jin *et al.*, 2007]. The result shown in Figure 1 supports the notion that atmospheric noise depends on the state of ENSO. That is, the El Niño SST anomaly allows favorable conditions for strong activity of the atmospheric transient zonal wind anomalies in the equatorial Pacific. Since there is a significant difference between noise that is independent of the ENSO state and noise that is dependent on ENSO state, as illustrated conceptual by Jin *et al.* [2007], the atmospheric transient variability may not only just trigger some ENSO events, but also impact ENSO growth rate and asymmetry. This argument will be discussed later in this study.

[7] The “atmospheric noise variance” in Figure 1 in fact includes a broad range of timescales from synoptic to the inter-seasonal timescale. To discern which frequency band of the atmospheric variability has a strong relation with ENSO, we first applied a Morlet wavelet analysis basis to a box-averaged unfiltered daily zonal wind anomaly. Then the 5-month moving-averaged wavelet power is linearly correlated with and regressed onto NINO3.4 SST with 12-month lead-lag span, as shown in Figure 2 for the three box-averaged zonal winds. For the wind anomalies averaged over 140 – 160°E , 5°S – 5°N , the wavelet power tends to lead the NINO3.4 SST for every frequency. This is consistent with the results from Figure 1. Significant correlations are found in the broad spectrum from 10 days to 90 days.

[8] For the box in the west of the dateline (160 – 180°E , 5°S – 5°N), the lead-lag relation is significantly different, depending on the frequency of the wind variability. For a

REG(NINO3.4 SST, Wavelet Power)




Figure 2. Lag-regression (contour) and correlation (shading) coefficients of wavelet power with respect to NINO3.4 SST. The wavelet power is obtained for the wind anomalies averaged over (a) 140–160°E, 5°S–5°N; (b) 160–180°E, 5°S–5°N; and (c) 160–180°W, 5°S–5°N. Lag (months) is indicated on the horizontal axes and period of wind variability (days) is indicated on the vertical axes. The unit of regression coefficients is $\text{m}^2 \text{s}^{-2} \text{K}^{-1}$.

shorter time scale (less than 20 days), the strong atmospheric noise variance leads El Niño by 3–12 months. Because the correlation is high even when the lag time is 12 months, the high-frequency variability can be a potential precursor of the ENSO prediction. Kug *et al.* [2005] showed that the low-frequency westerlies over the Western Pacific are observed prior to El Niño onset. The strong synoptic activity is possibly linked to the westerlies flow [Seiki and Takayabu, 2007a, 2007b]. However, the details of the dynamics behind this statistical relation are still not clear and will be pursued in further studies. Unlike the shorter time scale, the noise variance is simultaneously correlated to the NINO3.4 SST in the relatively longer time scale. The significant correlations are found in three frequency bands, which are the MJO frequency (20–60 days), extended MJO frequency range (60–90 days), and the inter-seasonal time scale (100–180 days). While this relation in MJO frequency is reported from several studies [Gutzler, 1991; Hendon *et al.*, 1999; Kessler, 2001; Tam and Lau, 2005], the relations in the two relatively lower frequency bands are not well documented, though Marcus *et al.* [2001] found that amplitude envelopes of the west equatorial Pacific wind stress in the extended MJO time scales significantly correlate with NINO3 SST at 8-month lag. The correlation in the period of 100–180 days is interesting; the correlation coefficients are robust with various moving windows (5 to 10 months). For the box in the east of the date line (160–180°W, Figure 2c), the correlation pattern is overall similar to that of Figure 2b. However, significant simultaneous correlations are found in the shorter time scales (5–20 days). From Figure 2, the fact that the relation between NINO3.4 SST and wind variance is strong for the variability in the frequency bands on intraseasonal timescale and beyond is noteworthy. The state-dependent atmospheric noises with a longer timescale

tend to have a more effective impact on ENSO because the noise-induced destabilization effect on ENSO growth rate is proportional to the lifetime scale of the state-dependent noise [Jin *et al.*, 2007].

[9] So far, we described the relationship between ENSO and atmospheric noise variances using the recent 26 years data (1980–2005). The characteristics of ENSO changed significantly in the late 1970s [Wang, 1995; An and Wang, 2000; An, 2004]. The amplitude and asymmetry of ENSO are enhanced in recent periods. It is possible that these changes of ENSO characteristics have been accompanied by change of the ENSO-noise relationship. To address this issue, the 15-years sliding regression and correlation between NINO3.4 SST and the noise variance are calculated as shown in Figure 3a. There is clearly a decadal modulation of the state-dependent noise. In general, the regression coefficients are gradually increasing over the central and eastern Pacific, even though relatively larger regression coefficients appeared at the early 1960s. On the other hand, the negative correlation is weakened over the 120–140°E during the recent period. As a result, the interaction between ENSO and the atmospheric noise is stronger in the recent period. Figure 3b shows the sliding regression and correlation between NINO3.4 SST and the wavelet power of the western box of the dateline (160–180°E) shown in Figure 2. These results indicate that the relation between ENSO and atmospheric noise is getting stronger.

[10] Jin *et al.* [2007] demonstrated that the state-dependent noise causes enhancing growth rate of El Niño and generates ENSO asymmetry. In their conceptual model, they assumed an ad hoc stochastic forcing as the form of $(1 + BT)\xi$ in the SST equation of the simple recharge-oscillator model [Jin, 1996, 1997; Burgers *et al.*, 2005]. Here, T represents normalized ENSO SST anomalies, such




Figure 3. 15-years sliding regression (contour) and correlation (shading) coefficients of (a) the noise variance and (b) the wavelet power of the western box of dateline (160–180°W, 5°S–5°N) with respect to the NINO3.4 SST. The period of wind variability (days) is indicated on the horizontal axis of Figure 3b. The unit of regression coefficients is $m^2 s^{-2} K^{-1}$.

as NINO3 SST, ξ is a red-noise process with constant variance and B is a non-dimensional parameter to represent simply the ENSO state-dependence of the noise forcing. Using the result in Figure 3, we may qualitatively estimate this parameter B by assuming a linear relationship between the variations in the noise variance and the NINO3.4 SST. The parameter B is thus proportional to $\frac{R}{\sigma_w^2 \sigma_N}$. Here, R is the regression coefficient (Figure 3) averaged in the central Pacific (160°E–140°W) and σ_w^2 is the mean variance of the atmospheric transient wind variability and σ_N is the standard deviation of NINO3.4 SST anomalies. As shown

in Figure 4, B is of relatively small value before the 1980s and increased significantly to a moderate value afterward.

[11] As it is noted by *Jin et al.* [2007], the impact of the state-dependent noise forcing on the stability of ENSO is proportional to B^2 . Because the value of B is nearly doubled before and after the middle 70s, the effect of the so-called noise-induced destabilization on the ENSO has quadrupled during this period. Thus the noise-induced destabilization effect could have played an important role in the enhancement of ENSO activity in the past few decades. This provides an alternative view to the observed




Figure 4. Changes of relative B value ($\frac{R}{\sigma_w^2 \sigma_N}$) with 15-year moving window. The detailed definition for relative B value is referred to in the text.

ENSO change, namely, the observed ENSO changes are generated by the slow background change [An and Jin, 2000; Fedorov and Philander, 2000], and internal ENSO dynamics [Timmermann and Jin, 2002].

4. Summary and Discussion

[12] The relationship between ENSO and atmospheric noise is examined using NCEP reanalysis data. We found that atmospheric noise variance over the central Pacific is simultaneously correlated to the NINO3.4 SST. That is, the atmospheric noise becomes significantly energetic during warm ENSO events. The significant relationship exists over most frequencies from the synoptic time scale to the inter-seasonal time scale. This observational study lends support to the notion that the atmospheric noise variability is modulated by the ENSO state. The extension of the warm pool and the low level westerlies give a favorable condition for the fast atmospheric variability during El Niño state [Seiki and Takayabu, 2007b]. When NINO3.4 SST is one degree above normal for moderate El Niño, the noise variance is changed by about 30–40 percent of the climatological noise variance over the central Pacific. In the case of strong El Niño, the change of the noise variance will be even larger than the climatological value. Recent theoretical work [Jin et al., 2007] proposed a new destabilization mechanism for ENSO through the so-called noise-induced instability due to state-dependent noise forcing. Our analysis suggests that this state-dependent noise has become stronger in the past few decades, which may contribute to the increases in the ENSO activity.

[13] **Acknowledgments.** This work is supported by the Korea Meteorological Administration Research and Development Program under Grant CATER_2007-4206 and the second stage of the Brain Korea 21 Project.

References

- An, S.-I. (2004), Interdecadal changes in the El Niño–La Niña asymmetry, *Geophys. Res. Lett.*, *31*, L23210, doi:10.1029/2004GL021699.
- An, S.-I., and F.-F. Jin (2000), An eigen analysis of the interdecadal changes in the structure and frequency of ENSO mode, *Geophys. Res. Lett.*, *27*, 2573–2576.
- An, S.-I., and B. Wang (2000), Interdecadal change in the structure of the ENSO mode and its impact on the ENSO frequency, *J. Clim.*, *13*, 2044–2055.
- Boullanger, J.-P., E. Durand, J.-P. Duvel, C. Menkes, P. Delecluse, M. Imbard, M. Lengaigne, G. Madec, and S. Masson (2001), Role of non-linear oceanic processes in the response to westerly wind events: New implications for the 1997 El Niño onset, *Geophys. Res. Lett.*, *28*, 1603–1606.
- Burgers, G., F.-F. Jin, and G. J. van Oldenborgh (2005), The simplest ENSO recharge oscillator, *Geophys. Res. Lett.*, *32*, L13706, doi:10.1029/2005GL022951.
- Duchon, C. (1979), Lanczos filtering in one and two dimensions, *J. Appl. Meteorol.*, *18*, 1016–1022.
- Eisenman, I., L. Yu, and E. Tziperman (2005), Westerly wind bursts: ENSO's tail rather than the dog?, *J. Clim.*, *18*, 5224–5238.
- Fedorov, A., and S. G. Philander (2000), Is El Niño changing?, *Science*, *288*, 1997–2002.
- Fink, A., and P. Speth (1997), Some potential forcing mechanisms of the year-to-year variability of the tropical convection and its intraseasonal (25–70 day) variability, *Int. J. Climatol.*, *17*, 1513–1534.
- Gutzler, D. S. (1991), Interannual fluctuations of intraseasonal variance of near-equatorial winds, *J. Geophys. Res.*, *96*, 3173–3185.
- Harrison, D. E., and G. A. Vecchi (1997), Westerly wind events on the tropical Pacific, 1986–95, *J. Clim.*, *10*, 3131–3156.
- Hendon, H., C. Zhang, and J. Glick (1999), Interannual variation of the Madden–Julian oscillation during austral summer, *J. Clim.*, *12*, 2538–2550.
- Jin, F.-F. (1996), Tropical ocean–atmosphere interaction, the Pacific cold tongue, and the El Niño–Southern Oscillation, *Science*, *274*, 76–78.
- Jin, F.-F. (1997), An equatorial ocean recharge paradigm for ENSO. Part I: Conceptual model, *J. Atmos. Sci.*, *54*, 811–829.
- Jin, F.-F., L. Lin, A. Timmermann, and J. Zhao (2007), Ensemble-mean dynamics of the ENSO recharge oscillator under state-dependent stochastic forcing, *Geophys. Res. Lett.*, *34*, L03807, doi:10.1029/2006GL027372.
- Kalnay, E., et al. (1996), The NCEP/NCAR 40-year reanalysis project, *Bull. Am. Meteorol. Soc.*, *77*, 437–471.
- Kessler, W. (2001), EOF representations of the Madden–Julian oscillation and its connection with ENSO, *J. Clim.*, *14*, 3055–3061.
- Kessler, W. S., and R. Kleeman (2000), Rectification of the Madden–Julian Oscillation into the ENSO cycle, *J. Clim.*, *13*, 3560–3575.
- Kessler, W. S., M. J. McPhaden, and K. M. Weickmann (1995), Forcing of intraseasonal Kelvin waves in the equatorial Pacific, *J. Geophys. Res.*, *100*, 10,613–10,631.
- Kirtman, B. P., K. Pegion, and S. Kinter (2005), Internal atmospheric dynamics and tropical Indo-Pacific climate variability, *J. Atmos. Sci.*, *62*, 2220–2233.
- Kug, J.-S., S.-I. An, F.-F. Jin, and I.-S. Kang (2005), Preconditions for El Niño and La Niña onsets and their relation to the Indian Ocean, *Geophys. Res. Lett.*, *32*, L05706, doi:10.1029/2004GL021674.
- Lau, K.-H., and N.-C. Lau (1992), The energetics and propagation dynamics of tropical summertime synoptic-scale disturbances, *Mon. Weather Rev.*, *120*, 2523–2539.
- Lengaigne, M., E. Guilyardi, J. P. Boulanger, C. Menkes, P. Delecluse, P. Inness, J. Cole, and J. Slingo (2004), Triggering of El Niño by westerly wind events in a coupled general circulation model, *Clim. Dyn.*, *23*, 601–620.
- Luther, D. S., D. E. Harrison, and R. A. Knox (1983), Zonal winds in the central equatorial Pacific and El Niño, *Science*, *222*, 327–330.
- Marcus, S. L., J. O. Dickey, and de Viron Olivier (2001), Links between intraseasonal (extended MJO) and ENSO timescales: Insights via geodetic and atmospheric analysis, *Geophys. Res. Lett.*, *28*, 3465–3468.
- McPhaden, M. J. (1999), Genesis and evolution of the 1997–1998 El Niño, *Science*, *283*, 950–954.
- Moore, A., and R. Kleeman (1999), The non-normal nature of El Niño and intraseasonal variability, *J. Clim.*, *12*, 2965–2982.
- Perez, C. L., A. M. Moore, J. Zavala-Garay, and R. Kleeman (2005), A comparison of the influence of additive and multiplicative stochastic forcing on a coupled model of ENSO, *J. Clim.*, *18*, 5066–5085.
- Seiki, A., and Y. N. Takayabu (2007a), Westerly wind bursts and their relationship with intraseasonal variations and ENSO. Part I: Statistics, *Mon. Weather Rev.*, *135*, 3325–3345.
- Seiki, A., and Y. N. Takayabu (2007b), Westerly wind bursts and their relationship with intraseasonal variations and ENSO. Part II: Energetics over the Western and Central Pacific, *Mon. Weather Rev.*, *135*, 3346–3361.
- Smith, T. M., and R. W. Reynolds (2004), Improved extended reconstruction of SST (1854–1997), *J. Clim.*, *17*, 2466–2477.
- Tam, C.-Y., and N.-C. Lau (2005), Modulation of the Madden–Julian Oscillation by ENSO: Inferences from observations and GCM simulations, *J. Meteorol. Soc. Jpn.*, *83*, 727–743.
- Timmermann, A., and F.-F. Jin (2002), A nonlinear mechanism for decadal El Niño amplitude changes, *Geophys. Res. Lett.*, *29*(1), 1003, doi:10.1029/2001GL013369.
- Vecchi, G. A., and D. E. Harrison (2000), Tropical Pacific sea surface temperature anomalies, El Niño and equatorial westerly events, *J. Clim.*, *13*, 1814–1830.
- Wang, B. (1995), Interdecadal changes in El Niño onset in the last four decades, *J. Clim.*, *8*, 267–285.
- Wittenberg, A. T. (2004), Extended wind stress analyses for ENSO, *J. Clim.*, *17*, 2526–2540.
- Wu, R., and B. P. Kirtman (2006), Changes in spread and predictability associated with ENSO in an Ensemble Coupled GCM, *J. Clim.*, *19*, 4378–4396.
- Yu, L., R. A. Weller, and W. T. Liu (2003), Case analysis of a role of ENSO in regulating the generation of westerly wind bursts in the western equatorial Pacific, *J. Geophys. Res.*, *108*(C4), 3128, doi:10.1029/2002JC001498.
- Zavala-Garay, J., C. Zhang, A. Moore, and R. Kleeman (2005), The linear response of ENSO to the Madden–Julian oscillation, *J. Clim.*, *18*, 2441–2459.
- Zhang, Q., and J. Gottschalk (2002), SST anomalies of ENSO and the Madden–Julian Oscillation in the Equatorial Pacific, *J. Clim.*, *15*, 2429–2445.

F.-F. Jin and J.-S. Kug, Department of Meteorology, School of Ocean and Earth Science and Technology, University of Hawaii at Manoa, 2525 Correa Road, HIG 350, Honolulu, HI 96822, USA. (jskug@hawaii.edu)

I.-S. Kang and K. P. Sooraj, Climate Environment System Research Center, Seoul National University, Sinlim-Dong, Kwanak-Gu, Seoul 151-742, Korea.